

What is the solar radiation value for ghana

What is the solar radiation value for ghana

ERA5-Land is a reanalysis dataset providing a consistent view of the volution of land variables over several decades at an enhanced resolution compared to ERA5. ERA5-Land has been produced by replaying the landcomponent of the ECMWF ERA5 climate reanalysis. Reanalysis combines modeldata with observations from across the world into a globally complete and consistent dataset using the laws of physics. Reanalysis produces data that goes several decades back in time, providing an accurate description of the climate of the past. This dataset includes all 50 variables as available onCDS.

The asset is a daily aggregate of ECMWF ERA5 Land hourly assets whichincludes both flow and non-flow bands. Flow bands are formed by collectingthe first hour's data of the following day which holds aggregated sum ofprevious day and while the non-flow bands are created by averaging allhourly data of the day. The flow bands are labeled with the "_sum"identifier, which approach is different from the daily data produced byCopernicus Climate Data Store, where flow bands are averagedtoo.

This problem is due to how the GRIB format saves data: it simplifies or "packs"the data into smaller, less precise numbers, which can introduce errors. These errors get worse when the data varies a lot.

Because of this, when we look at the data for a whole day to compute dailytotals, sometimes the highest amount of rainfall recorded at one time can seemlarger than the total rainfall measured for the entire day.

BandsNameUnitsDescriptiondewpoint_temperature_2mKTemperature to which the air, at 2 meters above the surface of theEarth, would have to be cooled for saturation to occur. It is ameasure of the humidity of the air. Combined with temperature and pressure, it can be used to calculate the relative humidity. 2m dewpoint temperature is calculated by interpolating between the lowestmodel level and the Earth's surface, taking account of the atmosphericconditions.

Temperature of air at 2m above the surface of land, sea or in-landwaters. 2m temperature is calculated by interpolating between thelowest model level and the Earth's surface, taking account of theatmospheric conditions.

Temperature of the surface of the Earth. The skin temperature is thetheoretical temperature that is required to satisfy the surface energybalance. It represents the temperature of the uppermost surface layer, which has no heat capacity and so can respond instantaneously tochanges in surface fluxes. Skin temperature is calculated differently over land and sea.

Temperature of the soil in layer 1 (0 - 7 cm) of the ECMWF IntegratedForecasting System. The surface is at 0 cm. Soil temperature is set atthe middle of each layer, and heat transfer is calculated at theinterfaces between



What is the solar radiation value for ghana

them. It is assumed that there is no heat transferout of the bottom of the lowest layer.

Temperature of water at the bottom of inland water bodies (lakes,reservoirs, rivers) and coastal waters. ECMWF implemented a lake modelin May 2015 to represent the water temperature and lake ice of all theworld's major inland water bodies in the Integrated ForecastingSystem. The model keeps lake depth and surface area (or fractionalcover) constant in time.

The thickness of ice on inland water bodies (lakes, reservoirs andrivers) and coastal waters. The ECMWF Integrated Forecasting System(IFS) represents the formation and melting of ice on inland waterbodies (lakes, reservoirs and rivers) and coastal water. A single icelayer is represented. This parameter is the thickness of that icelayer.

The temperature of the uppermost surface of ice on inland water bodies(lakes, reservoirs, rivers) and coastal waters. The ECMWF IntegratedForecasting System represents the formation and melting of ice onlakes. A single ice layer is represented.

This parameter describes the way that temperature changes with depthin the thermocline layer of inland water bodies (lakes, reservoirsand rivers) and coastal waters. It is used to calculate the lakebottom temperature and other lake-related parameters. The ECMWFIntegrated Forecasting System represents inland and coastal waterbodies with two layers in the vertical, the mixed layer above and thethermocline below where temperature changes with depth.

Contact us for free full report

Web: https://sumthingtasty.co.za/contact-us/ Email: energystorage2000@gmail.com WhatsApp: 8613816583346

